



## **Duke Orthopaedics: Upper Extremity Division**

*Christopher Klifto, MD & Katherine Batten, PA-C*

3609 SW Durham Dr, Durham, NC 27707

### **Discharge Instructions after Total Shoulder Arthroplasty or reverse Total Shoulder Arthroplasty (shoulder replacement)**

#### **Dressings:**

- A sling has been provided for you. Remain in your sling at all times. This includes sleeping in the sling. The sling may be removed for dressing and showering when cleared by physician.
- Leave the splint or dressing in place until your follow-up appointment
- The dressing **CANNOT** get wet.

#### **Pain Management/Medications:**

- Use ice (polar care unit) on the shoulder intermittently over the first 48 hours after surgery and as needed thereafter.
- Please resume all normal medications the day after surgery unless otherwise instructed.
- Please take a baby Aspirin (81mg) twice a day for 4 weeks to help prevent blood clot formation. If you take other blood thinners or are unable to take aspirin, please let us know.
- Pain medicine has been prescribed for you. Use your medication liberally as directed over the first 48 hours, and then begin to taper your use.
- No driving or drinking alcohol while taking narcotic pain medication
- You may take Extra Strength Tylenol or Tylenol in place or in conjunction with you prescribed pain medication. Do not exceed 4,000mg Tylenol per day (including Tylenol in certain prescribed pain medications like Percocet or Norco)

#### **Activity:**

- Avoid weight bearing with the operative upper extremity until your follow-up appointment. Do not push or pull with that arm. Do not lean on operative side through hand or elbow.
- No lifting greater than 1 pound: a cell phone, a small drink, or items of equivalent weight
- To prevent hand swelling: Move your fingers, squeeze stress ball, keep your hand elevated above your elbow, and keep your elbow above your heart
- Showering: sling may be removed, but keep arm at side. To wash operative armpit, lean forward and/or toward operative side so your arm passively moves away from the body. **DO NOT** raise your arm actively!

#### **Contact:**

- During business hours for urgent or routine questions, call Josh at the office: 919-403-3057
- After hours for emergent questions or concerns call and ask for the orthopaedic resident on call: 919-684-8111
- Contact the office or emergency on-call resident if you notice any of the following:
  - Excessive redness or warmth of the incisions
  - Drainage for more than 4 days after surgery
  - Fever of more than 101.5 F
  - Worsening numbness of the hand
  - Pain, redness, and swelling in your leg

## **Duke Orthopaedics: Upper Extremity Division**

*Christopher Klifto, MD & Katherine Batten, PA-C*

3609 SW Durham Dr, Durham, NC 27707

**-Go directly to the closest Emergency Room if you develop chest pain, difficulty breathing, or any complications that require immediate attention.**

Follow-up:

**You should see the Dr. Klifto or his PA Katie approximately 14 days after your surgery. A post-op appointment should have already been scheduled. If you don't have an appointment or you need to change the appointment, please call 919-471-9622.**